

Yanai T, Hirase S, Matsunoshita N, Yamamoto N, Ninchoji T, Kubokawa I, Mori T, Hayakawa A, Takeshima Y, Iijima K, Matsuo M.

Place of death of pediatric cancer patients in a single institute during 7 years.

Kobe J Med Sci. 58(2): E33-40, 2012.

Yamamoto T, Kawano S, Sugiyama D, Onishi T, Hayashi N, Takeshima Y, Kawai H, Hirata K, Matsuo M, Kumagai S.

Predicting scores for left ventricular dysfunction in Duchenne muscular dystrophy.

Pediatr Int. 54(3): 388-92, 2012.

Solyom S, Ewing AD, Hancks DC, Takeshima Y, Awano H, Matsuo M, Kazazian H H, Jr.

Pathogenic orphan transduction created by a nonreference LINE-1 retrotransposon.

Hum Mutat. 33(2): 369-71, 2012.

Shima Y, Nakanishi K, Hama T, Mukaiyama H, Togawa H, Hashimura Y, Kaito H, Sako M, Iijima K, Yoshikawa N.

Validity of the Oxford classification of IgA nephropathy in children.

Pediatr Nephrol. 27(5): 783-92, 2012.

Pham TV, Hartomo TB, Lee MJ, Hasegawa D, Ishida T, Kawasaki K, Kosaka Y, Yamamoto T, Morikawa S, Yamamoto N, Kubokawa I, Mori T, Yanai T, Hayakawa A, Takeshima Y, Iijima K, Matsuo M, Nishio H, Nishimura N.

Rab15 alternative splicing is altered in spheres of neuroblastoma cells.

Oncol Rep. 27(6): 2045-9, 2012.

Ohtsubo H, Morisada N, Kaito H, Nagatani K, Nakanishi K, Iijima K.

Alport-like glomerular basement membrane changes with renal-coloboma syndrome.

Pediatr Nephrol. 27(7): 1189-92, 2012.

Ogawa A, Kitamura S, Nakayama K, Sugiyama H, Morisada N, Iijima K, Makino H.

Right hypoplastic kidney.

Kidney Int. 82(9): 1037, 2012.

Nurputra DK, Nakagawa T, Takeshima Y, Harahap IS, Morikawa S, Sakaeda T, Lai PS, Matsuo M, Takaoka Y, Nishio H.

Paramyotonia congenita: from clinical diagnosis to in silico protein modeling analysis.

Pediatr Int. 54(5): 602-12, 2012.

Nishimura N, Hartomo TB, Pham TV, Lee MJ, Yamamoto T, Morikawa S, Hasegawa D, Takeda H, Kawasaki K, Kosaka Y, Yamamoto N, Kubokawa I, Mori T, Yanai T, Hayakawa A, Takeshima Y, Iijima K, Matsuo M, Nishio H.

Epigallocatechin gallate inhibits sphere formation of neuroblastoma BE(2)-C cells.

Environ Health Prev Med. 17(3): 246-51, 2012.

Manriquez ME, Makino A, Tanaka M, Abe Y, Yoshida H, Morioka I, Arakawa S, Takeshima Y, Iwata K, Takasaki J, Manabe T, Nakaya T, Nakamura S, Iglesias AL, Rossales RM, Mirabal EP, Ito T, Kitazawa T, Oka T, Yamashita M, Kudo K, Shinya K.

Emergence of HA mutants during influenza virus pneumonia.

Int J Clin Exp Pathol. 5(8): 787-95, 2012.

Malueka RG, Takaoka Y, Yagi M, Awano H, Lee T, Dwianingsih EK, Nishida A, Takeshima Y, Matsuo M.

Categorization of 77 dystrophin exons into 5 groups by a decision tree using indexes of plicing regulatory factors as decision markers.

BMC Genet. 13: 23, 2012.

Iijima K, Someya T, Ito S, Nozu K, Nakanishi K, Matsuoka K, Ohashi H, Nagata M, Kamei K, Sasaki S.

Focal segmental glomerulosclerosis in patients with complete deletion of one WT1 allele.

Pediatrics. 129(6): e1621-5, 2012.

Hashimoto F, Nishiumi S, Miyake O, Takeichi H, Chitose M, Ohtsubo H, Ishimori S, Ninchoji T, Hashimura Y, Kaito H, Morisada N, Morioka I, Fukuoka H, Yoshida M, Iijima K.

Metabolomics analysis of umbilical cord blood clarifies changes in saccharides associated with delivery method.

Early Hum Dev. 2012. Nov 21. [Epub ahead of print]

Harahap IS, Saito T, San LP, Sasaki N, Gunadi, Nurputra DK, Yusoff S, Yamamoto T, Morikawa S, Nishimura N, Lee MJ, Takeshima Y, Matsuo M, Nishio H.

Valproic acid increases SMN2 expression and modulates SF2/ASF and hnRNPA1 expression in SMA fibroblast cell lines.

Brain Dev. 34(3): 213-22, 2012.

Hama T, Nakanishi K, Shima Y, Mukaiyama H, Togawa H, Tanaka R, Hamahira K, Kaito H, Iijima K, Yoshikawa N.

Renal biopsy criterion in children with asymptomatic constant isolated proteinuria.

Nephrol Dial Transplant. 27(8): 3186-90, 2012.

Sonoyama A, Ebina Y, Morioka I, Tanimura K, Morizane M, Tairaku S, Minematsu T, Inoue N, Yamada H.

Low IgG avidity and ultrasound fetal abnormality predict congenital cytomegalovirus infection.

J Med Virol. 84(12): 1928-33, 2012.

Yamada H, Morizane M, Tanimura K, Morioka I, Ebina Y, Matsuda H, Yoshida M, Wakamatsu H, Furuya K, Murotsuki J, Saito S, Kawabata I, Takahashi Y, Ueda T, Sato Y, Kamitomo M, Maruyama Y, Hirano H, Yamada T, Minakami H, Yaegashi N, Minematsu T, Matsuoka R, Banzai M, Ikenoue T, for the Japanese congenital cytomegalovirus infection immunoglobulin fetal therapy study group.

A trial of immunoglobulin fetal therapy for symptomatic congenital cytomegalovirus infection.

J Reprod Immunol. 95(1): 73-9, 2012.

Morioka I, Morikawa S, Miwa A, Minami H, Yoshii K, Kugo M, Kitsunezuka Y, Enomoto M, Jikimoto T, Nakamura M, Yokoyama N, Nishio H, Matsuo M, Yamada H.

Culture-proven neonatal sepsis in Japanese neonatal care units in 2006-2008.

Neonatology. 102: 75-80, 2012.

Morioka I, Nonoyama S, Tanaka-Taya K, Ihara T, Sugaya N, Ueta I, Kumagai T, Okada K, Hosoya M, Okabe N, Morishima T.

Survey of Japanese infants younger than 3 months who were treated with oseltamivir for influenza: Safety of oseltamivir treatment.

Scand J Infect Dis. 44(8): 605-9, 2012.

Nakagawa T, Mure T, Yusoff S, Ono E, Harahap IS, Morikawa S, Morioka I, Takeshima Y, Nishio H, Matsuo M.

Acetaminophen administration in a Gilbert's syndrome patient.

Pediatr Int. 54(6): 934-6, 2012.

Miwa A, Morioka I, Yokota T, Shibata A, Matsuo K, Fujioka K, Koda T, Morikawa S, Nishio H, Yokoyama N, Nakamura H, Matsuo M, Yamada H.

Correlation and precision of serum free bilirubin concentrations measured by single and two peroxidase concentration methods in term or late-preterm newborn infants using a FDA-approved analyzer.

Clin Lab. 58: 507-14, 2012.

Fujioka K, Morioka I, Miwa A, Yokota T, Matsuo K, Morikawa S, Enomoto M, Shibata A, Morizane M, Yokoyama N, Yamada H, Matsuo M.

Renin is activated in monochorionic diamniotic twins with birthweight discordance who do not have twin-to-twin transfusion syndrome.

J Perinatol. 32: 514-9, 2012.

Sato Y, Morioka I, Miwa A, Yokota T, Matsuo K, Koda T, Fujioka K, Morikawa S, Shibata A, Yokoyama N, Takahashi K, Nishio H, Matsuo M.

Is the bilirubin/albumin ratio correlated with unbound bilirubin concentrations?

Pediatr Int. 54(1): 81-5, 2012.

- Mure T, Nakagawa T, Okizuka Y, Takami Y, Oyazato Y, Nagase H, Maruyama A, Adachi M, Takada S, Matsuo M.
Treatment of preterm infants with West syndrome: Differences due to etiology.
Pediatr Int. 54(6): 892-8, 2012.
- Thu Tran TH, Zhang Z, Yagi M, Lee T, Awano H, Nishida A, Okinaga T, Takeshima Y, Matsuo M.
Molecular characterization of an X(p21.2;q28) chromosomal inversion in a Duchenne muscular dystrophy patient with mental retardation reveals a novel long non-coding gene on Xq28.
J Hum Genet. 2012. Dec 6. [Epub ahead of print]
- Ota M, Takeshima Y, Nishida A, Awano H, Lee T, Yagi M, Matsuo M.
A G-to-T transversion at the splice acceptor site of dystrophin exon 14 shows multiple splicing outcomes that are not exemplified by transition mutations.
Genet Test Mol Biomarkers. 16(1): 3-8, 2012.
- Ito S, Kamei K, Ogura M, Udagawa T, Fujinaga S, Saito M, Sako M, Iijima K.
Survey of rituximab treatment for childhood-onset refractory nephrotic syndrome.
Pediatr Nephrol. 2012. Oct 10. [Epub ahead of print]
- Ishikura K, Yoshikawa N, Nakazato H, Sasaki S, Iijima K, Nakanishi K, Matsuyama T, Ito S, Yata N, Ando T, Honda M; Japanese Study Group of Renal Disease in Children.
Two-year follow-up of a prospective clinical trial of cyclosporine for frequently relapsing nephrotic syndrome in children.
Clin J Am Soc Nephrol. 7(10): 1576-83, 2012.
- Ikezumi Y, Honda M, Matsuyama T, Ishikura K, Hataya H, Yata N, Nagai T, Fujita N, Ito S, Iijima K, Kaneko T, Uemura O.
Establishment of a normal reference value for serum β 2 microglobulin in Japanese children: reevaluation of its clinical usefulness.
Clin Exp Nephrol. 2012. Jul 14. [Epub ahead of print]
- Uemura O, Honda M, Matsuyama T, Ishikura K, Hataya H, Nagai T, Ikezumi Y, Fujita N, Ito S, Iijima K.
Japanese Society for Pediatric Nephrology, the Committee of Measures for Pediatric CKD. Is the new Schwartz equation derived from serum creatinine and body length suitable for evaluation of renal function in Japanese children?
Eur J Pediatr. 171(9): 1401-4, 2012.
- Chikamoto H, Hattori M, Kuroda N, Kajiho Y, Matsumura H, Fujii H, Ishizuka K, Hisano M, Akioka Y, Nozu K, Kaito H, Shimizu M.
Pretransplantation combined therapy with plasmapheresis and rituximab in a second living-related kidney transplant pediatric recipient with a very high risk for focal segmental glomerulosclerosis recurrence.
Pediatr Transplant. (7): E286-90, 2012.

～英文論文 2012 年度～

Tsutsumi M, Kowa-Sugiyama H, Bolor H, Kogo H, Inagaki H, Ohye T, Yamada K, Taniguchi-Ikeda M, Toda T, Kurahashi H.

Screening of genes involved in chromosome segregation during meiosis I: in vitro gene transfer to mouse fetal oocytes.

J Hum Genet. 2012 Aug; 57(8): 515-22, 2012.