

## Corrigendum

### Corrigendum: Influences of Orogastric and Nasogastric Tubes on Sucking Pressure during Bottle Feeding in Immature Infants

TAKESHI KOMATSU<sup>1</sup>, TATSUYA FURUKAWA<sup>1,\*</sup>, DAISUKE KOBAYASHI<sup>2</sup>,  
SOTA IWATANI<sup>3</sup>, SEIJI YOSHIMOTO<sup>3</sup>, TOSHIHIKO YAMASHITA<sup>1</sup>,  
YUMI ENDO<sup>1</sup>, SAYAKA KATSUNUMA<sup>4</sup>, MASAHIDE OTSU<sup>4</sup>,  
and KEN-ICHI NIBU<sup>1</sup>

<sup>1</sup>*Department of Otolaryngology-Head and Neck Surgery, Kobe University Graduate School of Medicine, Kobe, Japan;*

<sup>2</sup>*Department of Pediatric Rehabilitation and Orthopedics, Hyogo Prefectural Kobe Children's Hospital, Kobe, Japan;*

<sup>3</sup>*Department of Neonatology, Hyogo Prefectural Kobe Children's Hospital, Kobe, Japan;*

<sup>4</sup>*Department of Otolaryngology, Hyogo Prefectural Kobe Children's Hospital, Kobe, Japan*

*\*Corresponding author*

Kobe J. Med. Sci., Vol. 69, No. 1, pp. E25-E32, 2023

In this article, the Ratio(post/pre) Data in Table IV were incorrect. This was due to an error during manuscript preparation, when creating Table IV, the values in Table III were erroneously adopted. Thus, the Ratio(post/pre) Data in Table IV has been updated. The P value Data in Table IV were correct, and these errors do not affect the results or conclusions of the article. We apologize for any confusion this may have caused. The original incorrect Ratio(post/pre) Data for Table IV are provided as Supplementary Information to this Corrigendum, for transparency. The original article has been corrected online.

## Supplementary Information

Table IV. Characteristics of ratio (post/pre)

Birth weight <1000 g group				
OG→NG (n = 3)	Ratio (post/pre)			p value
Sucking pressure	1.89	±	0.93	0.115
Area 1	2.94	±	2.28	0.014
Area 2	2.59	±	1.98	0.132
Intake	1.67	±	0.88	0.188
Birth weight ≥1000 g group				
OG→NG (n = 4)	Ratio (post/pre)			p value
Sucking pressure	1.06	±	0.21	0.558
Area 1	1.06	±	0.31	0.587
Area 2	0.98	±	0.10	0.935
Intake	1.14	±	0.70	0.616
NG→OFF (n = 3)	Ratio (post/pre)			p value
Sucking pressure	1.06	±	0.21	0.823
Area 1	1.06	±	0.31	0.361
Area 2	0.98	±	0.10	0.186
Intake	1.14	±	0.70	0.924

OG, orogastric tube; NG, nasogastric tube; OFF, oral feeding.

Values are expressed as mean ± standard error of the mean.  $P < 0.05$  was considered to indicate significance.